

REFERENCES

1. Lasslo, A., and P. D. Waller. 1957. 3-(*N,N*-Dialkylcarboxamido)piperidinoalkanes. *J. Org. Chem.* 22:837-838.
2. Lasslo, A., W. M. Marine, and P. D. Waller. 1956. Substituted piperidinecarboxamides. *J. Org. Chem.* 21:958-960.
3. Quintana, R. P., and W. A. Shrader. 1963. Synthesis of selected amides of mono- and bis(carboxypiperidino)alkanes. *J. Pharm. Sci.* 52:1186-1188.
4. Beasley, J. G., R. P. Quintana, and G. G. Nelms. 1964. The effect of piperidinecarboxamide derivatives on isolated human plasma cholinesterase. II. Variations in the amide function. *J. Med. Chem.* 7:698-701.
5. Quintana, R. P., T. D. Smith, and L. F. Lorenzen. 1965. Synthesis of carbamoylpiperidine-type cholinesterase inhibitors. *J. Pharm. Sci.* 54:785-787.
6. Quintana, R. P., and W. R. Smithfield. 1967. Substituted 1-benzyl-3-(*N,N*-diethylcarbamoyl)piperidine cholinesterase inhibitors. Relationships between molecular constitution, pK_a' values and partition coefficients. *J. Med. Chem.* 10:1178-1180.
7. Dachary-Prigent, J., J. Dufourcq, C. Lussan, and M. Boisseau. 1979. Propranolol, chlorpromazine and platelet membrane: a fluorescence study of the drug - membrane interaction. *Thromb. Res.* 14:15-22.
8. Vanderhoek, J., and M. B. Feinstein. 1979. Local anesthetics, chlorpromazine and propranolol inhibit stimulus-activation of phospholipase-A₂ in human platelets. *Mol. Pharmacol.* 16:171-180.
9. Chap, H. J., R. F. A. Zwaal, and L. L. M. Van Deenen. 1977. Action of highly purified phospholipases on blood platelets. Evidence for an asymmetric distribution of phospholipids in the surface membrane. *Biochim. Biophys. Acta.* 467:146-164.
10. Marcus, A. J., H. L. Ullman, and L. B. Safier. 1969. Lipid composition of subcellular particles of human blood platelets. *J. Lipid Res.* 10:108-114.
11. Quintana, R. P., A. Lasslo, M. Dugdale, L. L. Goodin, and E. F. Burkhardt. 1980. Effects of ethanol and of other factors on ADP-induced aggregation of human blood platelets in vitro. *Thromb. Res.* 20:405-415.
12. Quintana, R. P., A. Lasslo, G. S. Queen, and C. M. Baldwin. 1981. Aspirin analogs with enhanced interfacial activity. *J. Colloid Interface Sci.* In press.
13. Lasslo, A., and R. P. Quintana, Editors. 1973. Surface Chemistry and Dental Integuments. Charles C. Thomas, Springfield, IL.
14. Fisher, R. G., and R. P. Quintana. 1975. Surface-chemical studies on chlorhexidine and related compounds. II. Interactions with monomolecular-film systems. *J. Dent. Res.* 54:25-31.
15. Nys, G. G., and R. F. Rekker. 1974. The concept of hydrophobic fragmental constants (f-values). II. Extension of its applicability to the calculation of lipophilicities of aromatic and heteroaromatic structures. *Eur. J. Med. Chem.* 9:361-375.
16. Korolkovas, A. 1970. Essentials of Molecular Pharmacology; Background for Drug Design. Wiley-Interscience, New York; (a) 90, (b) 65-66.
17. Quintana, R. P. 1964. Relationships between the surface activity and cholinesterase inhibition of mono- and bis[3-(*N,N*-diethylcarbamoyl)piperidino]alkanes. *J. Pharm. Sci.* 53:1221-1223.
18. Quintana, R. P. 1965. Relationships between partition coefficients and cholinesterase inhibition of carbamoylpiperidinoalkanes. *J. Pharm. Sci.* 54:462-463.
19. Narahashi, T., and D. T. Frazier. 1971. Site of action and active form of local anesthetics. *Neurosci. Res.* 4:65-99.
20. Strichartz, G. R. 1973. The inhibition of sodium currents in myelinated nerve by quaternary derivatives of lidocaine. *J. Gen. Physiol.* 62:37-57.
21. Hille, B. 1977. Local anesthetics: hydrophilic and hydrophobic pathways for the drug-receptor reaction. *J. Gen. Physiol.* 69:497-515.
22. Quintana, R. P., A. Lasslo, M. Dugdale, and L. L. Goodin. 1981. Relationships between the chemical constitution of carbamoylpiperidines and related compounds, and their inhibition of ADP-induced human blood platelet aggregation. *Thromb. Res.* 22:665-680. Cf. Quintana, R. P., A. Lasslo and L. T. Greer. Interaction of human blood platelet aggregation inhibitors with phospholipid films. Submitted for publication.

ENVIRONMENTALLY INDUCED REVERSIBLE PROTEIN TRANSLOCATION ACROSS INNER MITOCHONDRIAL MEMBRANE

A. WAKSMAN

Centre de Neurochimie du Centre National de la Recherche Scientifique, 67084 Strasbourg Cedex, France

Four molecular mechanisms account for protein translocation across biological membranes: translocation via vectorial translation, via vectorial processing, via subunit dissociation, and via environmental induction. Whereas the first three mechanisms are vectorial, the last one is reversible (1). Fig. 1 describes the general methodology and observations that led us to the discovery of this last mechanism.

RESULTS AND DISCUSSION

Externalization of a protein such as aspartate aminotransferase (AAT) from the inner face of inner mitochondrial

membrane towards intermembranal fluid is shown in Fig. 1, step 1. This step is induced by movement effectors, e.g., succinate, fumarate, malate, pyruvate, citrate, phenylsuccinate, α -methylglutamate, etc., at concentrations ranging from 0 to 50 mM (2). Step 2 shows internalization of previously externalized AAT upon removal of the movement effector. This was demonstrated by testing the accessibility of the enzyme protein first on the outer face of the mitoplast (mitochondria from which the outer membrane was removed), then on the outer face of the corresponding inverted vesicle, which is equivalent to the inner face of the mitoplast (step 3). No permeant substrate, controlled




FIGURE 1 Environmentally induced aspartate aminotransferase movement within the mitochondria (10% AAT).

protease digestion, and specific antibody were used as criteria of accessibility. Outer membrane was separated from mitoplast after incubation of 10 mg mitochondrial protein with 1 mg digitonin (3). Specific reorganization of membrane and adjoining subcompartment following exposure to various movement effectors was estimated by showing: (a) the specific modification of intermembrane fluid protein patterns provoked by different movement effectors as revealed by polyacrylamide gel electrophoresis (2); and (b) the differential localization of AAT in the mitochondrial compartments in the presence of different movement effectors (3) (Table I). These results not only

TABLE I
INFLUENCE OF THE NATURE OF MOVEMENT
EFFECTOR ON THE SUBMITOCHONDRIAL
LOCALIZATION OF ASPARTATE AMINOTRANSFERASE

Movement effector	Percent of total aspartate aminotransferase activity in different submitochondrial compartments		
	Intermembrane space (fluid)	Inner membrane	Matrix
Sucrose	9	79	12
Phosphate	19	68	13
Acetate	20.5	72	7.5
Succinate	39	25	36
Fumarate	79	6	15
Oxaloacetate	77.5	11.5	11
Citrate	59.5	1.5	39

Results are expressed as percent of total aspartate aminotransferase activity. Total aspartate aminotransferase was constant to within $\pm 5\%$. Sucrose was 0.25 M. Effectors were at 2 $\mu\text{mol/mg}$ of mitochondrial protein.

confirm the specific action of effectors and repartition of membrane and perimembrane protein, but also demonstrate externalization. Inner membrane was separated from matrix after incubation of 1 mg mitoplast protein with 0.16 mg Lubrol WX.

This protein movement is dependent (4) on membrane fluidity and on temperature. The richer the membrane in polyunsaturated fatty acids, the lower the temperature of the break of the Arrhenius curve accounting for the movement effector induced protein externalization. In the case of the two populations of brain mitochondria (cell body and synaptosomal), when the ratio of saturated to polyunsaturated fatty acids varies from 1.07 to 0.62 the temperature of the break of the Arrhenius curve for AAT externalization varies from 27°C to 15°C.

Studies on the temperature-dependent movement of AAT in structurally intact mitochondria were performed by measuring reversible loss of AAT latency in the presence of 1.8 mM succinate using nonpermeant NADH at various temperatures (1, 3).

The results confirmed that AAT-reversible movements

TABLE II
DRIVING FORCE FOR PROTEIN MOVEMENT

Possible nature	Results	Conclusions
Energy-rich intermediate (ATP)	No inhibition of reversible movement by antimycin A, rotenone, oligomycin, 2-4 DNP, amytal, atractylate. Also triggered by phenylsuccinate, and α -methyl glutamate (2)	No energy rich link required
Modification of the primary structure of mobile protein	No modification of molecular weight of the "mobile AAT" as attested by gel electrophoresis on externalized, internalized and reexternalized ^{125}I enzyme*	No modification of primary structure required
Modulation of chemical potential of interacting constituents	"Spontaneous" internalization of mitochondrial aspartate aminotransferase within negatively charged liposomes as shown by latency of enzyme and inaccessibility to protease digestion (6)	Possible role of the modification of chemical potential

*Crémel, G., P. Hubert, and A. Waksman; data to be published.

occur *in situ* and showed that succinate externalization of AAT requires a more fluid membrane than for its internalization. As a matter of fact, internalization still occurs at temperatures below the thermotropic fusion point of the membrane for the externalization of the enzyme.

This asymmetrical recognition of both the externalizing signal and the molecule to be transitorily internalized was also suggested by cross-linking experiments. After treating mitochondria with dimethyladipimidate, externalization of AAT was no longer triggered by succinate, whereas in its absence, internalization of 70% of the enzyme still occurred on cross-linked organelles. This suggests that a greater rigidity or stability of the membrane could be required for the recognition and integration of protein signals as opposed to recognition of ionic signals (5). The driving force of this protein movement seems to be dependent on the chemical potential of both mobile protein and membrane as modulated by environment rather than by hydrolysis of an energy rich bond, or modification of the primary structure of the mobile protein (Table II).

Configurational energy and cognitive properties could thus be responsible for the externalization-internalization phenomenon.

Supported by grants from the Fondation par la Recherche Médicale Française n° 50.8131, and the Institut National de la Santé et de la Recherche Médicale No. 77.4.100.3 and 80.30.22.

Received for publication 13 April 1981.

REFERENCES

1. Waksman, A., P. Hubert, C. Crémel, A. Rendon, and C. Burgun. 1980. Translocation of proteins through biological membranes. A critical view. *Biochim. Biophys. Acta*. 604:209.
2. Waksman, A., and A. Rendon. 1974. Intramitochondrial intermembranal large amplitude protein movements. A possible novel aspect of membrane fluidity. *Biochimie*. 56:907.
3. Hubert, P., G. Crémel, A. Rendon, B. Sacko, and A. Waksman. 1979. Direct evidence for internalization of mitochondrial aspartate aminotransferase into mitoplasts. *Biochemistry*. 8:3119.
4. Crémel, G., G. Rebel, J. M. Warter, A. Rendon, and A. Waksman. 1976. Reversible intramitochondrial release of protein related to unsaturated fatty acids of membrane. *Arch. Biochem. Biophys.* 173:255.
5. Rendon, A., and A. Waksman. 1980. Translocation of mitochondrial aspartate aminotransferase through mitochondrial inner membrane. A cross-linking study with dimethyladipimidate. *Biochim. Biophys. Acta*. 603:178.
6. Furuya, E., Y. Yoshida, and K. Tagawa. 1979. Interaction of mitochondrial aspartate aminotransferase with negative charged lecithin liposomes. *J. Biochem. (Tokyo)*. 85:1157.

FORMATION OF THE ENVELOPE OF ROUS SARCOMA VIRUS AND VESICULAR STOMATITIS VIRUS FROM LOCALIZED LIPID REGIONS IN THE PLASMA MEMBRANE

GENEVA M. OMANN, JOSEPH CIMINO, JEFFREY E. PESSIN, AND MICHAEL GLASER
Department of Biochemistry, University of Illinois, Urbana, Illinois 61801 U.S.A.

The phospholipid polar head group composition of chicken embryo fibroblasts and enveloped viruses grown on these cells can be altered by supplementing the culture media with polar head group analogues. Studies of the incorporation of analogues into host cell plasma membranes and virus envelopes have shown that the viruses bud from localized lipid regions which are different from the average composition of the plasma membrane (1). This selective incorporation of phospholipids into the virus envelope may result from a specific requirement of the virus to regulate the envelope membrane structure. To test this hypothesis, the effects of altering the host cell plasma membrane lipid composition and fluidity on the viral envelope lipid composition and fluidity were determined. Measurements were made with 1,6-diphenyl-1,3,5-hexa-

triene (DPH) as a probe of membrane structure and fluidity using steady-state anisotropy methods and differential polarized phase fluorometry (2).

METHODS

Chicken embryo fibroblasts were cultured as previously described (3). RSV-T5-infected cells growing at 41°C were split into fresh medium, and half were incubated at 41°C and half at 36°C. After 32 h the medium was changed, and viruses and cells were harvested 12 h later. Polar head group modification of phospholipids was done on tertiary cultures using previously described methods (1). Cells were infected for 12 h with VSV (San Juan strain) and viruses and cells harvested after 36 h growth in choline- or *l*-2-amino-1-butanol-supplemented media. Virus purification, plasma membrane isolation, and phospholipid analysis were done as previously described (1). Membranes were labeled with DPH as previously described (4).

RESULTS

RSV-T5 is a Rous sarcoma virus mutant which infects cells and replicates equally well at 36°C and 41°C. At

Dr. Pessin's current address is the J. W. Wilson Laboratory, Brown University, Providence, Rhode Island 02912 U.S.A.